

Hemorrhoids

Did you know...


- Hemorrhoids are one of the most common ailments known.
- More than half the population will develop hemorrhoids, usually after age 30.
- Millions of Americans currently suffer from hemorrhoids.
- The average person suffers in silence for a long period before seeking medical care.
- Today's treatment methods make some types of hemorrhoid removal much less painful.

WHAT ARE HEMORRHOIDS?

Often described as "varicose veins of the anus and rectum", hemorrhoids are enlarged, bulging blood vessels in and about the anus and lower rectum. There are two types of hemorrhoids: external and internal, which refer to their location.

External (outside) hemorrhoids develop near the anus and are covered by very sensitive skin. If a blood clot develops in one of them, a painful swelling may occur. The external hemorrhoid feels like a hard, sensitive lump. It bleeds only if it ruptures.

Internal (inside) hemorrhoids develop within the anus beneath the lining. Painless bleeding and protrusion during bowel movements are the most common symptom. However, an internal hemorrhoid can cause severe pain if it is completely "prolapsed" - protrudes from the anal opening and cannot be pushed back inside.



WHAT CAUSES HEMORRHOIDS?

An exact cause is unknown; however, the upright posture of humans alone forces a great deal of pressure on the rectal veins, which sometimes causes them to bulge. Other contributing factors include:

- Aging
- Chronic constipation or diarrhea
- Pregnancy
- Heredity
- Faulty bowel function due to overuse of laxatives or enemas; straining during bowel movements
- Spending long periods of time (e.g., reading) on the toilet

Whatever the cause, the tissues supporting the veins stretch. As a result, the veins dilate; their walls become thin and bleed. If the stretching and pressure continue, the weakened veins protrude.

WHAT ARE THE SYMPTOMS?

If you notice any of the following, you could have hemorrhoids:

- Bleeding during bowel movements
- Protrusion during bowel movements
- Itching in the anal area
- Pain
- Sensitive lump(s)

DO HEMORRHOIDS LEAD TO CANCER?

No. There is no relationship between hemorrhoids and cancer. However, the symptoms of hemorrhoids, particularly bleeding, are similar to those of colorectal cancer and other diseases of the digestive system. Therefore, it is important that all symptoms are investigated by a physician specially trained in treating diseases of the colon and rectum. Do not rely on over-the-counter medications or other self-treatments. See a colorectal surgeon first so your symptoms can be properly evaluated and effective treatment prescribed.

HOW ARE HEMORRHOIDS TREATED?

Mild symptoms can be relieved frequently by increasing the amount of fiber (e.g., fruits, vegetables, breads and cereals) and fluids in the diet. Eliminating excessive straining reduces the pressure on hemorrhoids and helps prevent them from protruding. A sitz bath - sitting in plain warm water for about 10 minutes - can also provide some relief


With these measures, the pain and swelling of most symptomatic hemorrhoids will decrease in two to seven days, and the firm lump should recede within four to six weeks. In cases of severe, persistent pain, your physician may elect to remove the hemorrhoid containing the clot with a small incision. Performed under local anesthesia as an outpatient, this procedure generally provides relief.

Severe hemorrhoids may require special treatment, much of which can be performed on an outpatient basis.

- **Ligation** - the rubber band treatment - works effectively on internal hemorrhoids that protrude with bowel movements. A small rubber band is placed over the hemorrhoid, cutting off its blood supply. The hemorrhoid and

the band fall off in a few days and the wound usually heals in a week or two. This procedure sometimes produces mild discomfort and bleeding.

- **Injection and Coagulation** can also be used on bleeding hemorrhoids that do not protrude. Both methods are relatively painless and cause the hemorrhoid to shrivel up.
- **Hemorrhoidectomy** - surgery to remove the hemorrhoids - is the best method for the permanent removal of hemorrhoids. It is necessary when (1) clots repeatedly form in external hemorrhoids; (2) ligation fails to treat internal hemorrhoids; (3) the protruding hemorrhoid cannot be reduced; or (4) there is persistent bleeding. A hemorrhoidectomy removes excessive tissue that causes the bleeding and protrusion. It is done under anesthesia any may, depending upon circumstances, require hospitalization and a period of inactivity. Laser hemorrhoidectomies do not offer any advantage over standard operative techniques. They are also quite expensive, and contrary to popular belief, are no less painful.
- **Other treatments** include cryotherapy, BICAP coagulation and direct current. Cryotherapy, popular 20 years ago, consists of freezing hemorrhoidal tissue. It is not recommended for hemorrhoids because it is very painful. BICAP and direct current are methods that shrink the hemorrhoid. None of these treatments have gained widespread acceptance.



Rubber Band Ligation of Internal Hemorrhoids

- A. Bulging, bleeding, internal hemorrhoid
- B. Rubber band applied at the base of the hemorrhoid
- C. About 7 days later, the banded hemorrhoid has fallen off leaving a small scar at its base (arrow)

WHAT IS A COLON AND RECTAL SURGEON?

Colon and rectal surgeons are experts in the surgical and nonsurgical treatment of colon and rectal problems. They have completed advanced training in the treatment of colon and rectal problems in addition to full training in general surgery. Colon and rectal surgeons treat benign and malignant conditions, perform routine screening examinations and surgically treat problems when necessary.

The executive office of the 1,800-member American Society of Colon and Rectal Surgeons is located in the Chicago suburb of Arlington Heights. Board-certified colon and rectal surgeons complete a residency in general surgery, plus an additional year in colon and rectal surgery, and pass an intensive examination conducted by the American Board of Colon and Rectal Surgery.

For additional information or a [list of colorectal surgeons in your area](#), contact:

American Society of Colon and Rectal Surgeons
85 W. Algonquin Rd., Suite 550
Arlington Heights, IL 60005
Email: ascrs@fascrs.org